

# Study design and sampling intensity for demographic analyses of bear populations

Richard B. Harris, Charles C. Schwartz, Richard D. Mace,  
and Mark A. Haroldson

May 22, 2012





$\lambda$

How many bears – and of what sort – should we follow (and for how long) if our objective is to quantify  $\lambda$  with some specified level of precision?



## Demographic projection

- Age-specific vital rates
- Asymptotic
- Information-rich

## Empirical (C-MR)

- Population sizes (or estimates of seniority [Pradel approach])
- Real-time
- Information-limited

$\lambda$

## Demographic projection

- Age-specific vital rates
- Asymptotic
- Information-rich



“After spending  $x$  dollars monitoring the population for  $y$  years, and applying the most state-of-the-art demographic methods, we have determined that the population is either going up or it’s going down...”

$$\lambda = 1.02 (0.95 - 1.09)$$





- Denial



- Anger



- Bargaining



- Depression



- Acceptance



# How are bears typically studied?



Photo: Courtesy IGBST



Photo: Shannon Podruzny



How are bears typically studied?



Both photos: Steve Ard



# How many bears are typically followed annually?



Study Area/species	Females/yr	Litters/yr	Cubs/yr
Yellowstone (griz)	23.0	5.1	6.8
Montana (griz)	20.4	5.0	10.0
Alaska (griz)	22.8	7.4	11.7
B.C. (griz)	7.9	?	2.9
Alberta (black)	8.6	2.5	3.7
Arkansas (black)	31.1	15.4	3.5
Ontario (black)	35.9	11.9	22.8
Florida (black)	22.8	?	12.5

# How many years elapse before estimates of $\lambda$ are produced?

Study Area/species	Years elapsed	Source:
Yellowstone (griz)	19-20	Schwartz et al. 2006
Montana (griz)	6	Mace et al. 2012
Alaska (griz)	11	Kovach et al. 2006
B.C. (griz)	15	Hovey and McLellan 1996
Alberta (black)	6	Hebblewhite et al. 2003
Arkansas (black)	5-7	Clark and Eastridge 2006
Ontario (black)	11	Obbard and Howe 2008
Florida (black)	4-7	Hostetler et al. 2009

# How much uncertainty is typically present in vital rate estimates?

Study Area/species	Adult female survival	Cub survival	Fecundity
Yellowstone (griz)	1.4%*	5.0%	7.8%
Montana (griz)	2.2%	17.6%	12.3%
Alberta (black)	12.7%	16.7%	23.1%
Arkansas (black)	2.2%*	47.3%*	19.8%*

Values are: SE/Mean (%)

\* = Process variance only

# Doak et al. 2005

$$\begin{aligned} \text{Var}(\ln \hat{\lambda}_s) &= \sum_i \text{Var}(\hat{v}_i) \left( \frac{\partial \ln \lambda_s}{\partial v_i} \right)^2 \\ &+ \frac{1}{\hat{\lambda}_1^4} \sum \text{Var}(\hat{\sigma}_{i,E}) \left( \sum_j S_i S_j \rho_{i,j} \sigma_{j,E} \right)^2 \\ &+ \frac{1}{\hat{\lambda}_1^4} \sum \text{Var}(\hat{\rho}_{i,j}) (S_i S_j \sigma_{i,E} \sigma_{j,E})^2 \end{aligned} \quad (\text{Eq. A1})$$

*Ecology*, 86(5), 2005, pp. 1154–1163  
© 2005 by the Ecological Society of America

## UNDERSTANDING AND PREDICTING THE EFFECTS OF SPARSE DATA ON DEMOGRAPHIC ANALYSES

DANIEL F. DOAK,<sup>1,4</sup> KEVIN GROSS,<sup>2</sup> AND WILLIAM F. MORRIS<sup>3</sup>

<sup>1</sup>Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, California 95064 USA

<sup>2</sup>Department of Statistics, North Carolina State University, Raleigh, North Carolina 27695 USA

<sup>3</sup>Biology Department, Duke University, Box 90338, Durham, North Carolina 27708-0338 USA

**Abstract.** Demographic models are an increasingly important tool in population biology. However, these models, especially stochastic matrix models, are based upon a multitude of parameters that must usually be estimated with only a few years of data and limited

- Adult females monitored/yr: 10 – 60
- Litters monitored: 15% to 100% of females
- Years considered in estimates: 2 - 20

of demographic predictions: the sampling of small numbers of individuals within each year, so that vital rate estimates made for each transition period are uncertain; and the collection of data over only a small number of years, so estimates of the mean, variance,

of uncertainty around estimates of variance and correlation with limited data. Consequently, we ask: when is the cost of including estimates of variance and correlation in a demographic model worth the benefit? The basic idea behind this question is now well-publicized

# Growth and Sustainability of Black Bears at White River National Wildlife Refuge, Arkansas

JOSEPH D. CLARK<sup>a</sup>, United States Geological Survey, Southern Appalachian Research Branch, Lexington Science Center, University of Tennessee, Knoxville, TN 37996, USA

RIK EASTRIDGE, Arkansas Game and Fish Commission, Little Rock, AR 72201, USA

## Abstract

The black bear (*Ursus americanus*) population at White River National Wildlife Refuge is isolated and genetically distinct, but hunting occurs adjacent to refuge boundaries and females with cubs are removed annually for a reproductive census. We tagged and radio-collared bears to determine level of exploitation and compare methods for estimating population growth and sustainability. We captured 260 bears (133M:147F), 114 females, from 1988 through 2003. Survival estimates based on radio-collaring and mark-recapture indicated that hunting and translocations were significant sources of loss. Based on mark-recapture data (Pruett estimated), the annual population growth rate ( $\lambda$ ) averaged 1.00 (SE = 0.07) when translocation removals occurred and averaged 0.96 (SE = 0.18) when both harvest and translocation removals occurred. Estimates of  $\lambda$  based on a population simulation model (program HSPGM) averaged 1.00 (SE = 0.10) and 1.00 (SE = 0.11) when no removals occurred, 1.00 (SE = 0.00) and 1.00 (SE = 0.10) when translocation removals occurred, and 0.97 (SE = 0.05) and 1.00 (SE = 0.05) when both harvest and translocation removals occurred, depending on the survival rate estimates we used. The probability of population decline by >35% over a 10-year period ranged from 12.8 to 68.6%, given our estimated removal rates. We conclude that hunting and translocation losses are at or below the maximum the population is capable of sustaining. Although extinction risk of this isolated bear population are low over the near term, it should continue to be closely monitored by state and federal agencies. The mark-recapture method we used to estimate  $\lambda$  proved to be a reliable alternative to more costly population modeling methods. *UDJRNAL OF WILDLIFE MANAGEMENT* 76(4):1064–1071, 2009.

## Key words

abundance, black bear, population growth, Prue model, *Ursus americanus*, White River.

White River National Wildlife Refuge (NWR) in eastern Arkansas, USA, supports the largest remnant population of black bears in the Mississippi alluvial plain. Land-clearing for agriculture has caused losses of >80% of the bottomland hardwood habitat in the region (Neal 1990). The bear population at White River NWR was thought to have declined to about 25 animals by the 1940s (Dellinger 1942, Holder 1951) but has dramatically increased since that time (Smith 1973, Bowman et al. 1996, White 1996). Bears at White River NWR have a distinct genetic makeup, and issues regarding their taxonomic classification relative to the threatened Louisiana black bear have yet to be resolved (Calkins et al. 2003). Nuisance complaints have become more increasing bear population at White River Game and Fish Commission initiated a season on private lands adjacent to White address the nuisance bear issues and provide to the public. Additionally, we removed White River NWR for a reintroduction NWR, 160 km to the south (Weaver 2003). White River NWR bear population is managed because of its isolation, relatively value. Our objectives were to estimate the compare methods for estimating growth a bear population on the southern portion.

## Study Area

White River NWR is in the Lower Mississippi River (Bailey 1995) and encompasses

<sup>a</sup> E-mail: jclark@usgs.gov

1064

Desha, Monroe, and Phillips counties in eastern Arkansas. This long, narrow refuge varies from 5 to 16 km in width and contains a 145-km portion of the White River, just upstream from its confluence with the Mississippi River (Fig. 1). A network of ditches and bays connects more than 350 lakes within the refuge. White River NWR is approximately 65,000 ha in size, with roughly 2,000 ha of permanent water, 62,000 ha of forests, 360 ha of crops, and 400 ha of grassland. Agriculture is the primary use for lands surrounding the refuge, and so is the major crop followed by soybeans, corn, wheat, and sorghum. Winter and

The Journal of Wildlife Management 76(4):1064–1071, 2012. DOI:10.1002/jwm.1260

## Population Ecology

# Grizzly Bear Population Vital Rates and Trend in the Northern Continental Divide Ecosystem, Montana

RICHARD D. MACE<sup>a</sup>, Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, MT 59901, USA

DANIEL W. CARNEY, Blackfoot Tribe, Blackfoot Fish & Wildlife Department, Box 530, Browning, MT 59717, USA

TONYA CULBERTSON, Blackfoot Tribe, Blackfoot Fish & Wildlife Department, Box 530, Browning, MT 59717, USA

STACY A. COVILL, U.S. Geological Survey, Northern Rocky Mountain Science Center, Interagency Grizzly Bear Study Team, University of Montana, Missoula, MT 59717, USA

MARK A. HAROLDSON, U.S. Geological Survey, Northern Rocky Mountain Science Center, Interagency Grizzly Bear Study Team, University of Montana, Missoula, MT 59717, USA

RICHARD B. HARRIS, Department of Ecosystem and Conservation Science, College of Forestry and Conservation, University of Montana, Missoula, MT 59717, USA

JAMES JONKES, Montana Fish, Wildlife and Parks, 1201 Sprague Road, Missoula, MT 59904, USA

BRUCE MCLELLAN, British Columbia Ministry of Forest Research Branch, PO Box 1732, D'Arcy, BC, Canada V0N 1L0

MICHAEL MADEL, Montana Fish, Wildlife and Parks, Rocky Mountain Forest Field Office, Road 2002, Box 225, Glacier, MT 59942, USA

TIMOTHY L. MANLEY, Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, MT 59901, USA

CHARLES C. SCHWARTZ, U.S. Geological Survey, Northern Rocky Mountain Science Center, Interagency Grizzly Bear Study Team, University of Montana, Missoula, MT 59717, USA

CHRISTOPHER SERVEEN, U.S. Fish and Wildlife Service, University of Montana, Missoula, MT 59717, USA

GORDON STENHOUSE, Fish & Wildlife Research Station, Box 530, Browning, MT 59717, USA

JOHN S. WALLER, National Park Service, Glacier National Park, PO Box 124, West Glacier, MT 59936, USA

ERIK WENUM, Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, MT 59901, USA

**ABSTRACT.** We estimated grizzly bear (*Ursus arctos*) population vital rates and trend for the Northern Continental Divide Ecosystem (NCDE), Montana, between 2004 and 2009 by following radio-collared females and observing their fate and reproductive performance. Our estimates of dependent cub and yearling survival were 0.612 (95% CI: 0.309–0.818) and 0.682 (95% CI: 0.258–0.898). Our estimates of subadult and adult female survival were 0.852 (95% CI: 0.628–0.951) and 0.952 (95% CI: 0.892–0.980). From visual observations, we estimated a mean litter size of 2.00 cubs/litter. Accounting for cub mortality prior to the first observations of litters in spring, our adjusted mean litter size was 2.27 cubs/litter. We estimated the probabilities of females transitioning from one reproductive state to another between years. Using the stable state probability of 0.322 (95% CI: 0.262–0.382) for females with cub litters, our adjusted fecundity estimate ( $\lambda_0$ ) was 0.367 (95% CI: 0.273–0.461). Using our derived rates, we estimated that the population grew at a mean annual rate of approximately 3% ( $\lambda$ : 1.0306, 95% CI: 0.928–1.102), and 71.5% of 10,000 Monte Carlo simulations produced estimates of  $\lambda > 1.0$ . Our results indicate an increasing population trend of grizzly bears in the NCDE. Coupled with concurrent studies of population size, we estimate that over 1,000 grizzly bears reside in and adjacent to this recovery area. We suggest that monitoring of population trend and other vital rates using radio-collared females be continued. © 2011 The Wildlife Society.

**KEY WORDS.** grizzly bear, lambda, Montana, mortality, population trend, reproduction, survival, *Ursus arctos*.

The grizzly bear (*Ursus arctos*) occupies wilderness and non-wilderness habitats in the Northern Continental Divide Ecosystem (NCDE) of western Montana, USA. In 1975, this species was listed in the United States as Threatened

under the Endangered Species Act because of threats to habitat and populations. At that time, there was limited information on the ecological requirements of the species and its population status. The NCDE supports the largest population of 6 designated recovery areas for the species (U.S. Fish and Wildlife Service 1993) in the lower 48 states and is contiguous with populations to the north in Canada. Knowledge of population size and trend is useful in developing appropriate management programs for grizzly bears and to judge success of the recovery effort. In the Greater

Received: 9 November 2010; Accepted: 21 April 2011

Published: 6 December 2011

E-mail: rma@mt.gov

Muste et al. • NCDE Grizzly Population Trend



Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Biological Conservation 112 (2003) 415–425

BIOLOGICAL  
CONSERVATION

www.elsevier.com/locate/bcon

# Black bear (*Ursus americanus*) survival and demography in the Bow Valley of Banff National Park, Alberta

Mark Hebblewhite<sup>a,\*</sup>, Melanie Percy<sup>a</sup>, Robert Serrouya<sup>b</sup>

<sup>a</sup>Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E9

<sup>b</sup>R.S. - RR 1, Site 12, Campground 45, South Slave, BC, Canada V9G 2G9

Received 1 May 2002; received in revised form 6 October 2002; accepted 10 October 2002

## Abstract

We studied survival and demography of black bears, *Ursus americanus*, in Banff National Park (BNP) from 1994 to 2000 to test the efficacy of National Park protection. We monitored 25 radio-collared bears an average of 1.9 years each for a total of 51.8 bear-years. Eighty-two percent of all mortality ( $n=11$ ) was human-caused, composed of highway mortality (36%), management mortality (27%) and management relocation (18%). Survival was influenced by season and management status. Once bears became a management problem, survival (0.66) was lower than several subadult populations. Adult (0.84) survival was comparable to other unprotected or partly protected populations. Cub (0.68) and yearling (0.67) survival, and reproductive rate ( $\lambda$ ) = 0.47 female cub inter-birth interval) was slightly lower than other populations in western North America. We combined survival and reproductive rates in a preliminary post-birth pulse age-class Leslie matrix model and estimated population growth rate as 0.95 (95% simulated CI: 0.79–1.10). Sensitivity analyses showed  $\lambda$  was most sensitive to changes in adult female survival. Responsible management agencies should reduce adult female highway mortality and the likelihood of becoming a management problem, while continuing monitoring to reduce demographic analyses to adequately protect this population.

© 2002 Elsevier Science Ltd. All rights reserved.

**Keywords:** Black bear, Dens

## Introduction

Carnivores are extinction-prone taxa and extinction-prone taxa have distinctive characteristics (Noss et al., 1991; Al, 2001). Amongst these characteristics are: small range, low reproductive rate, and dependence on a single resource. Carnivores are also dependent on a single resource, and this dependence can be a source of risk (Noss et al., 1991; Noss et al., 1995).

Corresponding author.

E-mail address: mark.hebblewhite@ualberta.ca

0950-0687/\$ - see front matter © 2002 Elsevier Science Ltd.

PII: S0950-0687(02)00100-1

# Temporal, Spatial, and Environmental Influences on the Demographics of Grizzly Bears in the Greater Yellowstone Ecosystem

CHARLES C. SCHWARTZ<sup>a</sup>, U.S. Geological Survey, Northern Rocky Mountain Science Center, Interagency Grizzly Bear Study Team, Forestry Sciences Lab, Montana State University, Bozeman, MT 59717, USA

MARK A. HAROLDSON, U.S. Geological Survey, Northern Rocky Mountain Science Center, Interagency Grizzly Bear Study Team, Forestry Sciences Lab, Montana State University, Bozeman, MT 59717, USA

GARY C. WHITE, Department of Fishery and Wildlife Biology, Colorado State University, Fort Collins, CO 80523, USA

RICHARD B. HARRIS, Wildlife Biology Program, College of Forestry and Conservation, University of Montana, Missoula, MT 59812, USA

STEVE CHERRY, Department of Mathematical Sciences, Montana State University, Bozeman, MT 59717, USA

KIM A. KEATING, U.S. Geological Survey, Northern Rocky Mountain Science Center, Forestry Sciences Lab, Montana State University, Bozeman, MT 59717, USA

DAVE MOODY, Trophy Game Section, Wyoming Game and Fish Department, 260 Buena Vista, Lander, WY 82520, USA

CHRISTOPHER SERVEEN, U.S. Fish and Wildlife Service, University of Montana, Missoula, MT 59812, USA

## ABSTRACT

During the past 2 decades, the grizzly bear (*Ursus arctos*) population in the Greater Yellowstone Ecosystem (GYE) has increased in numbers and expanded in range. Understanding temporal, environmental, and spatial variables responsible for this change is useful in evaluating what likely influenced grizzly bear demographics in the GYE, and where future management efforts might benefit conservation and management. We used recent data from radio-collared bears to estimate reproduction (1985–2002) and survival (1985–2002) to evaluate combined effects on demographic vigor (lambda) ( $\lambda$ ). We explored the influence of an array of individual, temporal, and spatial covariates on demographic vigor.

We identified an important relationship between  $\lambda$  and where a bear resides within the GYE. This potential for a spatially-spatial dynamic in the GYE, coupled with concerns for managing available mortality, required our thinking about how management agencies might approach long-term conservation of the species. Consequently, we assessed the current spatial dynamics of the GYE grizzly bear population. Throughout, we followed the information-theoretic approach. We developed suites of a priori models that included individual, temporal, and spatial covariates that potentially affected reproduction and survival. We selected our best approximating models using Akaike's information criterion (AIC) and adjusted for small sample sizes and overdispersion (AIC<sub>c</sub> or QAIC<sub>c</sub>, respectively).

We provide recent estimates for reproductive parameters of grizzly bears based on 108 adult (>5 years old) females observed for 329 bear-years. We documented production of 104 litters with cub counts for 100 litters. Mean age of females producing their first litter was 5.81 years and ranged from 4 to 7 years. Proportion of subadult females that produced cubs at age 4–7 years was 9.8, 20.4, 36.4, and 100%, respectively. Mean (±SE) litter size ( $n=102$ ) was 2.0 ± 0.1. The proportion of litters of 1, 2, 3, and 4 cubs was 0.18, 0.61, 0.21, and 0.02, respectively. Mean yearling litter size ( $n=57$ ) was 2.0 ± 0.1. The proportion of litters containing 1, 2, 3, and 4 yearlings was 0.26, 0.51, 0.21, and 0.02, respectively. The proportion of adult females marked between years 1985 to 1990 was 0.05 to 0.08. The mean was 0.516 ± 0.03. Reproductive rate was estimated to be 0.518 female cub/year. We evaluated the probability of producing a litter of 2–3 cubs as well as to a suite of individual and temporal covariates using multinomial logistic regression. Our best models indicated that reproductive output, measured as cubs per litter, was most strongly influenced by indices of population size and white-tailed ptarmigan (Phasianidae) seed production. Our data suggest a possible density-dependent response in reproductive output, although parental mortality could have accounted for the correlation.

We analyzed survival of cubs and yearlings using radio-collaring of 46 unique female bears observed with 95 litters surviving 137 dependent years. We documented 42 deaths: 30 cubs, 5 yearlings, and 7 adults. Using a new survival estimator model (Program MARK), cub and yearling survival were most affected by winter weather in the GYE. Survival was highest for cubs and yearlings living outside Yellowstone National Park (YNP) but within the U.S. Fish and Wildlife Service (USFWS) Grizzly Bear Recovery Zone (RZ). Cubs and yearlings living inside YNP had lower survival rates, and those living outside the RZ had the lowest survival rates. Survival rates were negatively related to a population index, suggesting density dependence. Survival improved with higher white-tailed ptarmigan seed production, greater winter severity, larger female (mother's) age, and higher female (mother's) age. We tested theories of sexually selective interactions, but they were not supported.

We investigated factors influencing survival of adult and adult grizzly bears using data from 323 radio-collared bears monitored for 3,688 months. Telenary records were converted into monthly encounter histories, and survival was estimated using known fate data type in Program

E-mail: chuckschwartz@usgs.gov

Schwartz et al. • Demographics of the Yellowstone Grizzly




Lemons...







# GYE




# Hebblewhite black bears






# Lemonade...









## Acknowledgements:

- Dan Doak, Joe Clark, Mark Hebblewhite, John McDonald, Chris Servheen

### Citations:

Harris, R.B., C. C. Schwartz, M. A. Haroldson, and R.D. Mace. 2011. Study design and sampling intensity for demographic analyses of bear populations. *Ursus* 22: 24-36.

Doak, D.F. K. Gross, and W. F. Morris. 2005. Understanding and predicting the effects of sparse data on demographic analyses. *Ecology* 86(5): 1154–1163.